LECTURY CLASSIFICATION OF THIS PAGE					(
	KEPOKE DOCUI	MENTATION	PAGE			
18. REPORT SECURITY CLASSIFICATION	ELECTE	15 RESTRICTIVE				
AD A400 F04	P 1 9 1988	Distrib	ution Unlim	ited		
AD-A199 524	6) H 矣	5. MONITORING	ORGANIZATION R	REPORT NUMBER(5)	
64. NAME OF PERFORMING ORGANIZATION	6b. OFFICE SYMBOL (If applicable)	7a NAME OF MONITORING ORGANIZATION Office of Naval Research				
& ADDRESS (Gry, State and ZIP Code) Laboratory of Neuroscience, Bldg 8, Room 111 National Institutes of Healt		800 N.	ly. State. and ZIP Quincy Street on, VA 22217	et		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION	8b. OFFICE SYMBOL (If applicable)	}	T INSTRUMENT IC	ENTIFICATION N	JMBER	
Office of Naval Research	ONR	N00014-87				
8c. ADDRESS (City, State, and ZIP Code) 800 N. Quincy Street			FUNDING NUMBER			
Arlington, VA 22217-5000		PROGRAM ELEMENT NO. 61153N	PROJECT NO. RRO4108	TASK NO. 441 £726	WORK UNIT ACCESSION NO.	
11 TITLE (Include Security Classification) Neural Modulation of the Imm Ionophore Complex. 12 2875ONAL AUTHOR(S)	mune Response thro	ugh the Benz	odiazepine/(GABA Recepto	or Chloride	
Prince K. Arora, Ph.D.,	····				•	
13a. TYPE OF REPORT 13b. TIM ARTINIAI FROM	E COVERED 7-1-88	14. DATE OF REPO	ORT (Year, Month, St. 30. 1988	Day) 15. PAGE	COUNT 19	
16. SUPPLEMENTARY NOTATION			····			
77 COSATI CODES FIELD GROUP SUB-GROUP	18. SUBJECT TERMS (Benzodiaze)					
	NK activity	y; mitogen s	timulation;	MLR; I cell	s; Stress;	
19: ABSTRACT (Continue on reverse if necess			CNS Catorox	ic T Lymphoc	ute (CTL) (x)	
This project examines the complex (Supramolecular Computer Supramolecular Computer Suppression of allogeneic CTFG 7142 is dose-dependent, and the computer Section of FG 7142 are immune functions (CTL, NK, as a single dose of Alprazolam not "peripheral" benzodiazed receptors and the pathways control of immunity.	me role of the ben plex) in the contr IL response by the and that this supp nly, suggesting th cal Killer (NK) ce nd was still manif and MLR responses) (a triazolbenzodi pine receptors). I subserved by these	zodiazepine/ ol of immune benzodiazep ression is l at the FG 71 ll activity est 24 hr la was also ob azepine with hese results receptors m	functions. ine receptor ong lasting 42-induced was also su ter. A profe served 2 hr high affin suggest the	We have for inverse ag. FG 7142 suimmune supprepressed 2 hound suppressed after admirty for "centat the benze	and that the gonist, appressed ression may ar after ssion of mistration of ral" but addacepine	
20. DISTRIBUTION/AVAILABILITY OF ABSTRA		(ECURITY CLASSIFI U)			
Dr. J.A. Majde		225 TELEPHONE (202) 696	(Include Area Coo	ONR	JOENY	
	All other editions are o	obsolete.	DISTR	CLASSIFICATION ATTE MOTTUGE Lidurg rol bevo	PEMENT A	
	88 9 16	224		Autribution Unl		

ANNUAL REPORT OFFICE OF NAVAL RESEARCH CONTRACT N00014-87-G-0178

July 1, 1987 to June 30, 1988

Prince K. Arora and Phil Skolnick, Laboratory of Neuroscience, NIDDK, National Institutes of Health, Bethesda, MD 20892

I. INTRODUCTION

During the past two decades, it has been shown that the immune system can be modulated not only by "classical" means (Jerne, 1955; Jerne, 1974; Benacerraf and McDevitt, 1972; Gershon, 1974; McDevitt, 1980) but also through mechanisms controlled by the central nervous system (CNS) (Solomon, 1969; Besedovsky and Sorkin, 1977). For example, both psychosocial and environmental stressors have been shown to affect the humoral and cellular components of immunity in laboratory animals and man (Ader, 1981; Tekoma and Huey, 1985; Arora et al, 1987). Individuals with psychosis (Kovaleva et al, 1977), bereavement and depression (Barthrop et al, 1977), and emotional stress exhibit impaired immune reactivity (Solomon, 1969), and investigations of experimental animals subjected to stress by overcrowding or avoidance conditioning have shown impaired immune responsiveness (Rassmussen et al, 1959; Solomon, 1969).

laboratory has had a longstanding interest in the neurochemical bases of anxiety (Tallman et al, 1980; Skolnick and Paul, 1983). Pharmacological, biochemical and behavioral evidence suggests that the benzodiazepine/GABA receptor chloride ionophore complex ("supramolecular complex") mediates the anti-anxiety effects of benzodiazepines, barbiturates, and pharmacologically important agents (Usdon et al, 1982). Several lines of evidence suggest that the supramolecular complex is involved in the physiological control of stress and anxiety (Ninan et al, 1982; Havoundjian et al, 1987). Since the role of the "supramolecular complex" in the neural modulation of immunity had not been investigated, we initiated such studies. Thus, we recently found that the administration of the benzodiazepine receptor (BzR) "inverse agonists" FG 7142 (N'-methyl-ß-carboline-3-carboxamide) and DMCM (3-carbomethoxy-4-ethyl-6,7-dimethoxy-ßcarboline) produced a profound suppression of T-cell functions in mice (Arora et al, 1987). Since \(\beta\)-carbolines like FG 7142 have been demonstrated to produce a BzR-mediated behavioral, somatic, and endocrine syndrome reminiscent of stress or anxiety in rodents and primates, including man (Ninan et al, 1982; Dorow et al, 1983; Insel et al, 1984), our findings (Arora et al, 1987) suggest that the benzodiazepine receptors in the CNS and the pathways subserved by these receptors may be important in the neural control of immunity.

II. PROGRESS REPORT

- A. IMMUNOMODULATION THROUGH THE BENZODIAZEPINE/GABA RECEPTOR CHLORIDE IONOPHORE COMPLEX (SUPRAMOLECULAR COMPLEX) IN THE CNS:
 - 1. Cytotoxic T-lymphocyte (CTL) Response Studies:
- a). Dose-Kinetics of FG 7142-Induced Immunosuppression: The immunoregulatory effects of FG 7142 were studied in vivo by administering varying doses of FG 7142 to groups of NFR/N mice (Small Animal Section, NIH, Behtesda, MD). FG 7142 (12.5-100 mg/kg, i.p.) was injected in 0.15 ml of vehicle (20% Emulphor). The placebo group received an equal volume of vehicle. Spleens were removed 24 hr later and the CTL response measured as described (Arora and Shearer, 1981). Results presented in Fig. 1 indicate that these doses of Fg 7142 significantly suppressed the CTL response. Furthermore as the dose of Fg 7142 was increased, greater suppression of the CTL response resulted, with 25 mg/kg being the optimal dose. These results suggested that suppression of the CTL response by FG 7142 is dose-dependent.
- FG-7142-Induced Immunosuppression is long-lasting: In preliminary studies, we had shown that 24 h after administration B-carbolines, both mitogen stimulated T-cell of proliferation and allogeneic CTL response were suppressed (Arora 1987). The length and magnitude of suppression are unknown. Animals were administered with 25 mg/kg of FG 7142, and at different time periods, animals were sacrificed and the CTL response measured as described (Arora and Shearer, 1981). Suppression of the CTL response was manifest even after 24 days suggesting that the suppression of the CTL response by FG 7142 is very long lived (Fig. 2). It would be of interest to extend this time course study to determine how long this suppression by FG 7142 lasts.
- c). Influence of Gender on Stress-induced Immunosuppression: Both male and female NFR/N mice were administered with FG 7142 (25 mg/kg) and 24 h later animals were sacrificed and the allogeneic CTL response measured. The placebo animals received vehicle only. Results as shown in Fig. 3 indicate that FG 7142 allogeneic CTL response in male mice only. Administration of doses even several-fold higher (100 mg/kg) failed to produce a similar effect in female mice. These results suggest that FG 7142-induced immunosuppression may be sexually dimorphic. It would be of interest to investigate the mechanisms r through which females resistant to stress-induced appear immunosuppression. Such studies would include an examination of the dose-effect relationship, duration of immunosuppression, and determination of suppression on specific T-cell populations in both sexes.

2. Natural Killer (NK) Cell Activity Studies:

During the first year, we also investigated whether the supramolecular complex modulates another immune parameter, NK safor



A-1

activity. Male B10.BR mice (Jackson Laboratories, Harbor, ME) were injected with FG 7142 (5-50 mg/kg) or an equal volume of vehicle. Spleens were removed 2 and 24 hr later and NK cell activity measured using chromium-51 (5 Cr) release assay as described (Arora and Shearer, 1981; Petitto et al, 1988). A dosedependent supression of NK cell activity was observed both at 2 hr (Fig. 4A) and 24 hr (Fig. 4B) after administration of FG 7142. A similar dose-dependent suppression of NK cell activity was observed at other effector:target (E:T) cell ratios (100:1 and 25:1) (data not shown). The doses of FG 7142 needed to suppress NK cell activity (Petitto et al, 1988) were consistent with those that produce both behavioral and endocrine changes in rodents reminiscent of stress or anxiety (File and Pellow, 1985; Stephens and Kehr, 1985) and those that suppressed T-cell functions (Arora et al, 1987). Pretreatment of mice with a specific, high affinity 15-1788 (10 mg/kg) 15 min prior to antagonist Ro administration of FG 7142 (25 mg/kg) resulted in a significant reduction of this suppression (Fig. 5). In this series of FG 7142 suppressed NK cell activity by 35.6% experiments, (compared with vehicle treated animals) which was reduced to 16.6% in mice pretreated with Ro 15-1788 (Fig. 5). Ro 15-1788 did not reduce NK cell activity when administered alone (Fig. 5).

Several observations in this study suggest that suppression of NK cell activity by FG 7142 is mediated via occupation of BzR in the CNS. Direct addition of FG 7142 (1 $\mu M-$ 10 μM) to the 51Cr release assays during a four hr incubation period had no effect on NK cell activity (data not shown). Furthermore, neither Ro 15-1788 nor inverse agonist FG 7142 bind with high affinity to peripheral benzodiazepine receptors (pBzR) (Marangos et al, 1982; Schoemaker et al, 1983) that are present on cells of the immune system (Zavala et al, 1985; Ruff et al, 1985; Moingeon et al, 1985; Zavala and Lenfant, 1987). Finally, the antagonism of FG 7142-induced suppression of NK cell activity by Ro 15-1788 is consistent with the ability of this compound to the effects of both BzR agonists (i.e. substances with benzodiazepine-like qualities) and inverse agonists (Skolnick and 1983). These findings suggests that the BzR agonists may be useful tools to study neural-immune interactions, and support the hypothesis (Arora et al, 1987) that the pathways subserved by the "supramolecular complex" may play an important role in the neural modulation of immunity.

Recent studies have demonstrated that the Long-Sleep (LS) and Short-Sleep (SS) mouse lines, bidirectionallly selected for their hypnotic sensitivities to a single dose of ethanol, are also differentially sensitive to other depressants such as barbiturates (McIntyre and Alpern, 1985, 1986; Marley et al, 1986) and benzodiazepines (McIntyre and Alpern, 1986), as well as convulsants such as 3-carbomethoxy-\$\beta\$-carboline, picrotoxin, and bicuculline (Philips and Dudek, 1983; McIntyre and Alpern, 1986). Thus, LS and SS mouse lines represent a unique genetic model which can be utilized to assess the role of the supramolecular complex in the neural modulation of immune functions. Since the

well-described difference in drug sensitivities of these lines through inherent differences appears to be mediated biochemical and biophysical properties of the supramolecular (Marley and Wehner, 1986; McIntyre et al, 1988), the complex assessment of NK cell function could be accomplished without confounding pharmacological intervention. Spleen cells from male LS and SS mice (Institute for Behavioral Genetics, University of Colorado, Boulder, CO), were tested for NK cell activity by using a 51Cr release assay as described (Arora and Shearer, 1981; Petitto et al, 1988). NK cytotoxic activity ranged from 6.0-16.9% in the LS mice and 3.6-7.0% in SS mice, respectively (Fig. 6). The NK cell activity of the LS line was higher than the SS line at each E:T ratio tested, with differences ranging from 67-142%. [Significant differences in the total numbers of cells per spleen were also observed between these lines. The number of viable cell per spleen was 68% higher in the LS line (178.8 \pm 15.3 x 10 $^{\circ}$) than in the SS line (106.3 \pm 6.5 \times 10°) (p<.001, Student's ttest)]. Since NK cell activity is assayed with equal number of effector spleen cells from each line, the greater number of splenic leukocytes in LS mice, thus, greatly enhanced the genetic differences in NK cell activity between LS and SS. with observations, in concert the findings benzodiazepine ligands affect immune functions (Arora et al, 1987), provide additional support for the hypothesis that the "supramolecular complex" (in the CNS) regulates NK cell activity.

3). Effect of Alprazolam on Selected Aspects of Immunity:

THE PARTY OF THE PROPERTY OF THE PARTY OF TH

Previous studies have demonstrated that anxiolytic benzodiazepines can modulate immune function (Descotes et al, 1982; Okimura & Nagata 1986; Pericic et al, 1987: Zavala & These effects have generally been attributed to Lenfant 1987). modulation of "peripheral" rather than "central" benzodiazepine receptors (Zavala & Lenfant 1987). "Peripheral" benzodiazepine receptors have been identified on components of the immune system such as macrophages (Zavala & Lenfant 1987) and human monocytes et al. 1985). In order to determine whether immune modulation by benzodiazepine receptor agonists is mediated via "central" benzodiazepine (i.e. the benzodiazepine/GABA receptor chloride channel complex), we examined in this study the effects of alprazolam (a triazolobenzodiazepine with high affinity for "central" but not "peripheral" benzodiazepine receptors) selected aspects of cellular immunity.

A single dose of alprazolam (0.5 or 1.0 mg/kg, i.p.) was administered to male B10.BR mice. At different time intervals (2, 2.5 or 24 hr later), selected aspects of immune function were examined. A profound suppression of immune function was observed two hr after injection. This suppression was manifest as a decrease in mitogen-stimulated T and B lymphocyte proliferation. A significant reduction in mixed leucocyte reaction (MLR) and allogeneic cytotoxic T lymphocyte (CTL) response was also observed upon administration of alprazolam. However, the immunosuppression produced under these conditions appeared short

lived. At 2.5 hr, only the CTL and MLR responses were suppressed whereas 24 hr after the initial dose of alprazolam, none of the parameters were different from those measures in vehicle treated mice (Table I).

These data suggest that benzodiazepine anxiolytic drugs may have significant though short-lived effects on immune function through activation of "central" receptors, since alprazolam has high affinity for "central" benzodiazepine receptors but very low affinity for the "peripheral" receptors. However, in order to assess the implication of these data for clinical use of antianxiety agents in the human population, we will have to delineate further the dose range (0.05-5.0 mg/kg) and time course of the immunosuppressive effects of alprazolam.

Arora et al (1987) have recently reported a profound inhibition of immune function after administration of the anxiogenic \(\beta\)-carbolines FG 7142 and DMCM. These anxiogenic drugs act as antagonists to benzodiazepines at the benzodiazepine/GABA receptor chloride channel complex (Bruun-Meyer 1987). Hence in future experiments we will study a possible interactive effect between such anxiogenic agents and alprazolam. By carefully examining the time interval required to see an interactive effect between administration of the \(\beta\)-carbolines and alprazolam, we will be able to relate activation of the receptor complex to the effects on the immune system.

III. REFERENCES :

- 1. Ader, R. (1981) (Ed.), Psychoneuroimmunology (Academic Press, Orlando).
- 2. Arora, P.K; and Shearer, G.M. (1981). Non-MHC-linked genetic control of murine cytotoxic T-lymphocyte responses to hapten-modified syngeneic cells. J. Immunol. 127: 1822-1828.
- 3. Arora, P.K; Hanna, E.E; Paul, S.M; and Skolnick, P. (1987). Suppression of the immune response by benzodiazepine receptor inverse agonists. J. Neuroimmunol. 15: 1-9.
- 4. Bartrop, R.W; Lazarus, L; Luckherst, E; Kiloh, L.G; and 'Penny, R. (1977). Depressed lymphocyte function after bereavement. Lancet. 1: 834-836.
- 5. Benacerraf, B; and McDevitt H.O. (1972). Histocompatibility-linked immune response genes. Science. 175: 273-279.
- 6. Besedovsky, H. O; and Sorkin, E. (1977). Network of immune-neuroendocrine interactions. Clin. exp. Immunol. 27: 1-12.
- 7. Bruun-Meyer S.E. Progr. Neuro-Psychopharmacol. & Biol. Psychiat. 11: 365-387, 1987.
- 8. Descotes J., R. Tedone and J.C. Evreux. Immun. Letters 5: 41-

- 43, 1982.
- 9. Dorow, R; Horowski, R; Paschelke, G; Amin, M; and Braestrup, C. (1983). Severe anxiety induced by FG 7142, a ß-carboline ligand for benzodiazepine receptors. Lancet ii: 98-99.
- 10. File, S; and Pellow, S. (1985). The effects of putative anxiogenic compounds, FG 7142, CGS 8216, and Ro 15-1788 on the rat corticosterone response. Physiol. Behav. 35: 587-90.
- 11. Gershon, R.K. (1974). T-cell control of antibody production. Contemp. Top. Immunobiol. 3: 1-25.
- 12. Havoundjian, H; Trullas, R; Paul, S; and Skolnick, P. (1987). A physiological role of the benzodiazepine receptor complex. In "Molecular Mechanisms of Neuronal Responsiveness" (Ehrlich, Y; Kornecki, E; Berry, W and Lenox, R. eds,), Plenum Press, New York.
- 13. Insel, T; Ninan, P; Aloi, J; Jimerson, D; Skolnick, P; and Paul, S. (1984). A benzodiazepine receptor mediated model of anxiety: Studies in non-human primates and clinical implications. Arch. Gen. Psych. 41: 741-750.
- 14. Jerne, N.K. (1955). The natural selection theory of antibody formation. Proc. Natl. Acad. Sci. 41: 849.
- 15. Jerne, N.K. (1974). The immune system: A web of v-domains. The Harvey Lectures. Academic Press, New York.
- 16. Kovaleva E.S; Bonartsev P.D; and Prilipko L.L. (1977). Response of lymphocytes of patients with schizophrenia to phytomitogens Concanavalin A and Phytohemagglutinin. Bull. exp. Biol. Med. 84: 1136-1139.
- 17. Marangos, P; Patel, J; Boulenger, J.P; and Clark-Rosenberg, R. (1982). Characterization of peripheral-type benzodiazepine binding sites in brain using [3H]Ro 5-4864. Mol. Pharmacol. 22, 26-32.
- 18. Marley, R.J; Miner, L.L; Wehner J.M; and Collins, A.C. .(1986). Differential effects of central nervous system depressants in long-sleep and short-sleep mice. J. Pharmacol. and Exp. Ther. 238: 1028-1033.
- 19. Marley, R.J; and Wehner, J.M. (1986). GABA enhancement of flunitrazepam binding in mice selectively bred for differential sensitivity to ethanol. Alcohal Drug Res. 7: 25-32.
- 20. McDevitt, H.O. (1980). Regulation of the immune response by the Major Histocompatibility Complex System. N. Engl. J. Med. 303: 1514-1517.

- 21. McIntyre T.D; and Alpern, H.P. (1985). Reinterpretation of the literature indicates differential sensitivities of long-sleep and short-sleep mouse lines are not specific to alcohal. Psychopharmacology 87: 379-389.
- 22. McIntyre T.D; and Alpern, H.P. (1986). Thiopental, phenobarbital, and chlordiazepoxide induce the same differences in narcotic reaction as ethanol in long-sleep and short-sleep selectively-bred mice. Pharmacol. Biochem. and Behav. 24: 895-898.
- 23. McIntyre, T.D; Trullas, R; and Skolnick, P. (1988).
 Differences in the biophysical properties of the
 benzodiazepine/GABA receptor chloride channel complex in
 Long Sleep (LS) and Short Sleep (SS) mouse lines. J.
 Neurochem. 51: 642-647.
- 24. Moingeon, P.H; Bidart, J.M; Alberici, C; and Bohuon, C. (1983). Characterization of a peripheral-type benzodiazepine binding site on human circulating lymphocytes. Eur. J. Pharmacol. 92: 147-149.
- 25. Ninan, P; Insel, T; Cohen, R; Cook, J; Skolnick, P; and Paul, S. (1982). Benzodiazepine receptor-mediated experimental "Anxiety" in Primates. Science 218: 1332-1334.
- 26. Okimura T. and I. Nagata. J. Immunopharmac. 8: 327-346, 1986.
- 27. Pericic D., H. Manev, M. Boranic, M. Poljak-Blazi and N. Lakic. NY Acad. Sci. 496: 450-458, 1987.
- 28. Petitto, J.P; Skolnick, P; and Arora, P.K. (1988). Suppression of Natural killer cell activity by the benzodiazepine receptor inverse agonist, FG 7142. Brain, Behavior and Immunity. (submitted).

SANDON COORDE NECESCO CONSIDER SPANOR DISABLE NATIONAL PROCESSE NATIONAL

- 29. Phillips T.J; and Dudek, B.J. (1983). Bicuculline-induced seizures in mice which differ in sensitivity to ethanol. Neurosci. Abstr. 9: 1240.
- 30. Rassmussen A. F; Spencer E. S; and Marsh J. T. (1959).

 Decrease in susceptibility of mice to Passive Anaphylaxis
 following Avoidance-learning stress. Proc. Soc. exp. Biol.

 Med. 100: 878-879.
- 31. Ruff, M.R; Pert, C.B; Weber, R.J; Wahl, S.M; and Paul, S.M. (1985). Benzodiazepine receptor mediated chemotaxis of human monocytes. Science 229: 1281-1282.
- 32. Schoemaker, H; Boles, R; Horst, W.D; and Yamamura, H. (1983). Specific high-affinity binding sites for [3H]Ro 5-4864 in rat brain and kidney. J. Pharmacol. Exp. Ther. 225:61-69.

- 33. Skolnick, P; and Paul, S.M. (1983). New concepts in the neurobiology of anxiety. J. Clin. Psychiat. 44: 12-19.
- 34. Solomon, G.F. (1969). Emotions, Stress, The central nervous system and Immunity. Int. Ann. N. Y. Acad. Sci. 164: 335-337.
- 35. Stephens, D; and Kehr, W. (1985). B-carbolines can enhance or antagonize the effects of punishment in mice. Psychopharm. 85: 143-147.
- 36. Tallman, J; Paul, S; Skolnick, P; and Gallager, D. (1980). Receptors for the age of anxiety: Molecular pharmacology of the benzodiazepines. Science. 207: 274-281.
- 37. Tecoma, E; and Huey, L. (1985). Psychic distress and the Immune Response. Life Sci. 36: 1799-1812.
- 38. Usdon, E; Skolnick, P; Tallman, J; Greenblat, D; and Paul, S. eds. (1982). In "Pharmacology of Benzodiazepines". McMillan Press, Ltd. London.
- 39. Zavala, F; Haumont, J; and Lenfant, M. (1985). Interaction of benzodiazepines with mouse macrophages. Eur. J. Pharmacol 106: 561-566.
- 40. Zavala, F; and Lenfant, M. (1987). Peripheral benzodiazepines enhance the respiratory burst of macrophage-like P338D1 cells stimulated by arachadonic acid. Int. J. Immunopharmac. 9/3: 269-274.
- 41. Zavala F. and M. Lenfant. (1987). Benzodiazepines and PK 11195 exert immunomodulating activities by binding on a specific receptor on macrophages. NY Acad. Sci 496: 240-249.

IV. PUBLICATIONS (Year 1):

Manuscripts:

- 1. Arora, P.K.: (1988). Neuromodulation of Natural Killer (NK) Cells. In "Neuroimmunomodulation" (Ed. Edward J. Goetzl) Alan R. Liss, Inc. (submitted)
- 2. Petitto, J.M., Skolnick, P., and Arora, P.K.: (1988). Suppression of Natural Killer (NK) Cell Activity by FG 7142, A Benzodiazepine Receptor "Inverse Agonist". Brain Behavior and Immunity (submitted)
- 3. Petitto, J.M., McIntire, T.D., Skolnick, P., and Arora, P.K.: (1988). Natural Killer (NK) Cell Activity in the Long-Sleep (LS) and Short-Sleep (SS) Mouse Lines: A Possible Association Between Alcohal Sensitivity and Cellular Immunity.

 J. Neuroimmunology. (submitted)
- 4. Arora, P.K., Fride, E., Petitto, J., Waggie, K., and Skolnick.: (1988). Are Morphine-Induced Alterations in Immune Function a Predisposing Factor for AIDS.

 Nature (to be submitted)
- 5. Ostrowski, N., Kress, D.W., Hagan, A., and Arora, P.K. (1988). Sexual Behavior Suppresses Immune Function.

 Nature (submitted)
- 6. Ostrowski, N., Kress, D.W., Hagan, A., and Arora, P.K. (1988). Sexual Behavior Suppresses Antibody Production in the Golden Hamster (Mesocricetus auratus). Brain, Behavior and Immunity (submitted)
- 7. Kress, D.W., Ostrowski, N.L., and Arora, P.K.: (1988). Mating Suppresses Splenic Natural Killer (NK) Cell Cytotoxicity in Male Golden Hamsters
 Brain, Behavior and Immunity. (submitted)

Abstracts:

en a benedica de la compacta del compacta del compacta de la compacta del la compacta de la comp

- 1. Petitto, J.M., Skolnick., and Arora, P.K.: (1988). Suppression of Natural Killer Cell Activity by FG 7142, A Benzodiazepine Receptor "Inverse Agonist". FASEB Summer Research Conference on "Neuroimmunomodulation". Copper Mountain, Colorado.
- 2. Fride, E., Skolnick, P., and Arora, P.K.: (1988). Transient Immunosuppressive Effects of Alprazolam in Mice. FASEB Summer Research Conference on "Neuroimmunomodulation". Copper Mountain, Colorado.
- 3. Fride, E., Collins, R.L., and Arora, P.K.: (1988). Different Immune Competence and "Anxiety Level" in Lateralized Mice. Society for Neuroscience Meeting, Toronto, Canada.

4. Arora, P.K., Fride, E., Petitto, J., Waggie, K., and Skolnick, P.: (1988). Morphine-Induced Modulation of the Immune System: Implications for AIDS. IV International Conference on AIDS. Stockholm, Sweden.

TRAINING ACTIVITIES: Two post-docs (Ester Fride, Ph.D., and John Petitto, M.D.) and two summer students (Douglas Kress and Bradford McRae) have been trained during the first year of the project.

AWARDS AND FELLOWSHIPS:

- 1). Mentor, NIH Summer Student Fellowship to Dr. Arora.
- 2). Mentor, The Armenian Assembly Summer Intern Program. Armenian Assembly of America, Washington, D.C.
- 3). International Travel Award to Dr. Arora to attend IV International Conference on AIDS, Stockholm, Sweden, 1988.

Figure. 1

DOSE-KINETICS OF FG 7142-INDUCED SUPPRESSION ON THE CTL RESPONSE




Figure. 2

TIME-KINETICS OF FG 7142-INDUCED SUPPRESSION ON THE CTL RESPONSE







Figure. 3










TABLE I Effects of alprazolam (ALP) on different parameters of immune function at 2, 2.5 or 24 hours after administration to male B10.BR mice.

Treatment	CTL (%1ysis)	MLR (cpm)	Conc A (cpm)	PHA (cpm)	LPS (cpm)
2 HOURS					
CONTROL	55±24	43661±3496	40980±5423	9286±1493	4958±328
ALP. 0.53	37±5²	28190±1872²	27884±3363	5288± 9932	3972±113
ALP. 1.0	28±8²	23804±1327²	22837±6000	6984± 7181	4080±442
2.5 HOURS					
CONTROL	62±2	41502±2034	50926±6960	13885±1683	75381±6757
ALP. 1.0	40±52	27969±40262	61251±6219	12724±1782	71456±3938
24 HOURS					
CONTROL	45±3	68625±8090	46350±2346	15465±2120	65880±6977
ALP. 1.0	51±4	66351±4856	43218±1230	12223±1155	62217±3579
1) p<0.05 2) p<0.01 3) mg/kg 4) sem Conc-A=concanava LPS=lipopolysacc	1) p<0.05 2) p<0.01 3) mg/kg 4) sem Conc-A=concanavalin-A LPS=lipopolysaccharide	0.1U/culture; 10U/culture; n=5	fo	PHA=phytohemagglutinin r each group.	2.5U/culture;

DISTRIBUTION LIST

Behavioral Immunology Program

Annual, Final and Technical Reports (one copy each except as noted)

INVESTIGATORS

Dr. Itamar B. Abrass
Department of Medicine
University of Washington
Harborview Medical Center
Seattle, WA 98104

Dr. Prince K. Arora NIDDK, Bldg. 8, Rm. 111 National Institutes of Health Bethesda, MD 20892

Dr. Andrew S. Baum
Department of Medical Psychology
Uniformed Services University
of Health Sciences, B3050
4301 Jones Bridge Road
Bethesda, MD 20814-4799

Dr. Charles A. Bowles Merrifield Research Lab, Inc. P.O. Box 2362 Merrifield, VA 22116-2362

Dr. Karen Bulloch Helicon Foundation 4622 Sante Fe Street San Diego, CA 92109

Dr. Michael D. Cahalan Department of Physiology and Biophysics University of California, Irving Irvine, CA 92717

Dr. Donald A. Chambers Health Sciences Center University of Illinois at Chicago P.O. Box 6998 Chicago, IL 60680

Dr. Christopher L. Coe Department of Psychology Harlow Primate Laboratory University of Wisconsin Madison, IL 53715

Dr. Sheldon Cohen
Department of Psychology
Carnegie-Mellon University
Pittsburgh, PA 15213

Dr. Walla L. Dempsey
Dept. of Microbiology and
Immunology
The Medical College of
Pennsylvania
3300 Henry Avenue
Philadelphia, PA 19129

Dr. David L. Felten
Department of Anatomy
University of Rochester
School of Medicine
601 Elmwood Avenue
Rochester, NY 14642

Dr. John F. Hansbrough Department of Surgery UCSD Medical Center 225 Dickinson Street San Diego, CA 92103

Dr. Robert L. Hunter
Department of Pathology
Emory Univ. School of
Medicine
WMB 760
Atlanta, GA 30322

Dr. Terry C. Johnson Division of Biology Ackert Hall Kansas State University Manhattan, KS 66506

Dr. Sandra Levy University of Pittsburgh School of Medicine 3811 O'Hara Street Pittsburgh, PA 15213

Dr. Lester Luborsky
Department of Psychiatry
308 Piersol Building/Gl
University of Pennsylvania Hospital
Philadelphia, PA 19104

Dr. Steven F. Maier Department of Psychology University of Colorado Campus Box 345 Boulder, CO 80309

Dr. Diana S. Malcolm
Department of Surgery, USUHS
Uniformed Services University
of Health Sciences
4301 Jones Bridge Road
Bethesda, MD 20814-4799

Dr. Michael H. Melner Department of Reproductive Biology Oregon Regional Primate Center 505 N.W. 185th Avenue Beaverton, OR 97006

Dr. Vera B. Morhenn
Department of Dermatology
Stanford University Medical School
Stanford, CA 94305

Dr. Jose R. Perez-Polo Gail Borden Bldg., Rm. 436 University of Texas Medical Branch Galveston, TX 77550-2777

Dr. Howard R. Petty Department of Biological Sciences Wayne State University Detroit, MI 48202

Dr. Bruce S. Rabin Clinical Immunopathology Childrens Hospital University of Pittsburgh School of Medicine Pittsburgh, PA 15213

Dr. Seymour Reichlin Director, Clinical Study Unit New England Medical Center Hospitals, Inc. 171 Harrison Avenue Boston, MA 02111 Dr. Eric M. Smith
Department of Psychiatry
University of Texas
Medical Branch
Galveston, TX 77550

Dr. Ross R. Vickers, Jr. Naval Health Research Ctr. Bldg. 346 P.O. Box 85122 San Diego, CA 92138

Annual, Final and Technical Reports (one copy each except as noted)

endere de la contraction de la

ADMINISTRATORS

Dr. Jeannine A. Majde, Code II4ISB (2 coples) Scientific Officer, Immunology Program Office of Naval Research 800 N. Quincy Street Arlington, VA 22217-5000

Administrator (2 copies) (Enclose DTIC Form 50) Defense Technical Information Center Building 5, Cameron Station Alexandria, VA 22314

Administrative Contracting Officer ONR Resident Representative (address varies - obtain from business office)

Annual and Final Reports Only (one copy each)

Program Manager Biological/Human Factors Division Office of Naval Research, Code 125 800 N. Quincy Street Arlington, VA 22217-5000

Program Manager Support Technology Directorate Office of Naval Technology, Code 223 800 N. Quincy Street Arlington, VA 22217-5000

DoD ACTIVITIES

Commanding Officer Naval Medical Center Washington, DC 20372

Commanding Officer
Naval Medical Research & Development Command
National Naval Medical Center
Bethesda, MD 20814

Director, Infectious Diseases Program Center Naval Medical Research Institute National Naval Medical Center Bethesda, MD 20814

Commander

Chemical and Biological Sciences Division Army Research Office, P.O. Box 12211 Research Triangle Park, NC 27709

Commander

U.S. Army Research and Development Command Attn: SGRD-PLA Fort Detrick Frederick, MD 21701

Final and Technical Reports Only

Director, Naval Research Laboratory (6 copies) Attn: Technical Information Division, Code 2627 Washington, DC 20375 Commander
USAMRIID
Fort Detrick
Frederick, MD 21701

Directorate of Life Sciences Air Force Office of Scientific Researc Bolling Air Force Base Washington, DC 20332

Library
Armed Forces Radiation Research
Institute
Bethesda, MD 20814-5145